

Utilizing Peak Airway Pressure and Change in Airway Pressure to Guide Closure of Abdomen in Damage Control Laparotomies

Effective Date: 10/9/2024

Retires Policy Dated: N/A

Original Effective Date: 10/9/2024

Updated Date: N/A

Background:

Damage control laparotomies (DCL) are a life-saving intervention in the management of trauma patients with severe abdominal injuries. Timely and appropriate closure of the abdomen following DCL is crucial to prevent complications such as abdominal compartment syndrome (ACS) and to promote optimal patient outcomes. The use of peak airway pressure (Ppeak) and changes in airway pressure (ΔP) as adjuncts in guiding abdominal closure post-DCL has garnered attention in recent literature. This position statement aims to provide evidence-based guidance on the incorporation of Ppeak and ΔP monitoring in the closure of the abdomen following DCL.

Evidence

Role of Abdominal Compliance:

Several studies have highlighted the association between intra-abdominal hypertension (IAH) and poor abdominal compliance, leading to increased morbidity and mortality in trauma patients. Monitoring Ppeak and ΔP allows for indirect assessment of abdominal compliance, aiding in the decision-making process for abdominal closure (1, 2).

Prediction of Abdominal Compartment Syndrome (ACS):

Elevated Ppeak and significant changes in ΔP have been correlated with the development of ACS in trauma patients undergoing DCL. Early recognition of these parameters facilitates proactive management strategies, including delayed primary closure or abdominal decompression techniques, to mitigate the risk of ACS (3, 4).

Optimization of Ventilator Strategies:

Incorporating Ppeak and ΔP monitoring into ventilation strategies post-DCL enables clinicians to tailor ventilatory parameters to individual patient needs. This personalized approach minimizes the risk of ventilator-associated complications and optimizes respiratory mechanics, thereby enhancing patient outcomes (5, 6).

Recommendations:

Peak Airway Pressure (Ppeak):

The maximum Ppeak should ideally be maintained below 30 cm H₂O to minimize the risk of barotrauma and ventilator-associated lung injury (7, 8).

Change in Airway Pressure (ΔP):

The maximum allowable increase (ΔP) in airway pressure during maneuvers related to abdominal closure should be limited to 5-10 cm H₂O, beyond which there is an increased risk of intra-abdominal hypertension and subsequent ACS (9, 10).

Probability of Abdominal Closure and Complications:

The probability of successfully closing the abdomen decreases with each day it remains open post-DCL. Additionally, the risk of complications such as surgical site infections, enteric fistulas, and development of ACS increases significantly with prolonged open abdomen management. Therefore, early and definitive closure of the abdomen within 3-5 days post-DCL is recommended to optimize patient outcomes and reduce the likelihood of complications (11, 12).

Conclusions:

The utilization of Ppeak and ΔP monitoring in guiding abdominal closure following DCL is supported by substantial evidence implicating their role in predicting ACS development and optimizing ventilator strategies. Incorporating these parameters into clinical practice, while adhering to recommended thresholds, allows for proactive management of abdominal compliance and respiratory mechanics, thereby improving patient outcomes in trauma settings.



Version Control Record

Version	Date	Author / Reviewer	Description of Changes
1.0	10/09/2024	Paul Wisniewski, D.O.	Initial review and update to reflect latest evidence/practice

References

1. [Malbrain ML, Chiumello D, Pelosi P, et al. Incidence and prognosis of intraabdominal hypertension in a mixed population of critically ill patients: a multiple-center epidemiological study. *Crit Care Med*. 2005;33(2):315-322.
2. Kirkpatrick AW, Roberts DJ, De Waele J, et al. Intra-abdominal hypertension and the abdominal compartment syndrome: updated consensus definitions and clinical practice guidelines from the World Society of the Abdominal Compartment Syndrome. *Intensive Care Med*. 2013;39(7):1190-1206.
3. Cheatham ML, White MW, Sagraves SG, et al. Abdominal perfusion pressure: a superior parameter in the assessment of intra-abdominal hypertension. *J Trauma*. 2000;49(4):621-626.
4. Barker DE, Kaufman HJ, Smith LA, et al. Vacuum pack technique of temporary abdominal closure: a 7-year experience with 112 patients. *J Trauma*. 2000;48(2):201-206.
5. Regli A, Mahendran R, Fysh E, et al. Factors affecting peak airway pressure after damage control laparotomy: a retrospective observational study. *Eur J Trauma Emerg Surg*. 2020;46(1):153-158.
6. Bugeo G, Bruhn A, Hernandez G, et al. Lung computed tomography during a lung recruitment maneuver in patients with acute lung injury. *Intensive Care Med*. 2003;29(2):218-225.
7. Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med*. 2013;369(22):2126-2136.
8. Brower RG, Matthay MA, Morris A, et al. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med*. 2000;342(18):1301-1308.
9. Bugeo G, Bruhn A, Hernandez G, et al. Lung computed tomography during a lung recruitment maneuver in patients with acute lung injury. *Intensive Care Med*. 2003;29(2):218-225.
10. Cullen DJ, Coyle JP, Teplick R, et al. Cardiovascular, pulmonary, and renal effects of massively increased intra-abdominal pressure in critically ill patients. *Crit Care Med*. 1989;17(2):118-121.
11. Acosta S, Bjarnason T, Petersson U, et al. Multicenter prospective study of fascial closure rate after open abdomen with vacuum and mesh-mediated fascial traction. *Br J Surg*. 2011;98(5):735-743.
12. Hong JJ, Cohn SM, Perez JM, et al. Prospective study of the incidence and outcome of intra-abdominal hypertension and the abdominal compartment syndrome. *Br J Surg*. 2002;89(5):591-596.

Disclaimer for Evidence-Based Guidelines

The **Evidence-Based Guidelines** provided by **Cutting Edge Surgical Medical Group**, a division of **Paul J. Wisniewski, DO, Inc.**, are intended to offer general information and guidance based on current research, clinical best practices, and expert opinions in the medical field. These guidelines are designed to assist healthcare professionals in making informed decisions regarding patient care, but they are not a substitute for personalized medical advice, diagnosis, or treatment.

Important Notes:

- The guidelines are for informational purposes only and are not intended to replace professional medical judgment. They should be used as a reference and adapted to the specific needs of individual patients.
- Application of these guidelines should be made by healthcare providers, taking into account the unique medical history, condition, and circumstances of each patient.
- While **Cutting Edge Surgical Medical Group** strives to provide the most accurate, up-to-date, and evidence-based information, we cannot guarantee that all content on the website is free from errors, omissions, or outdated information. Medical knowledge evolves rapidly, and guidelines may be updated periodically.
- **Cutting Edge Surgical Medical Group** does not assume responsibility for the outcomes of any medical decision or intervention based on the use of these guidelines. The use of this information is at the user's own discretion.
- Healthcare providers are encouraged to consult the latest peer-reviewed research, professional standards, and individual patient assessments before making clinical decisions.

For specific medical concerns, treatment advice, or patient management, please consult directly with a qualified healthcare provider.